of symptoms and erosions or ulceration associated with GERD, the recommended dosage is 20 mg twice daily.

ADVERSE EFFECT:
Reducid is generally well tolerated, however headache, dizziness, GI disturbances and fatigue have rarely been reported.

STORAGE:
Store below 30° C in a dry place, product from light. To be dispensed on the prescription of a registered medical practitioner only. Keep out of the reach of children.

PRESENTATION:
Reducid suspension is available in 120ml Bottle.
Reducid 20mg tablets are available in Blister pack of 20's.
Reducid 40mg tablets are available in Blister pack of 20's.

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر خشک جگہ پر رکھیں،
روشنی سے بچائیں۔ صرف رجسٹرڈ ڈاکٹر کے نسخے پر فروخت کریں۔
بچوں کی پہنچ سے دُور رکھیں۔

Manufactured by:
Platinum
Pharmaceuticals (Pvt) Ltd.
A-20, North Western Industrial Zone,
Bin Qasim, Karachi-75020, Pakistan.

Mfg. Lic No. 000415

Reducid
(Famotidine)

رڈیوسِڈ
(فیموٹیڈین)

Suspension 10mg / 5ml
Tablets 20 & 40mg

COMPOSITION:
Reducid Suspension 10mg / 5ml:
Each 5ml Contains:
Famotidine ..... 10 mg
Product Complies Platinum Specs

Reducid Tablets 20mg:
Each Film Coated Tablets Contains:
Famotidine ..... 20 mg
Product Complies U.S.P. Specs

Reducid Tablets 40mg:
Each Film Coated Tablets Contains:
Famotidine ..... 40 mg
Product Complies U.S.P. Specs

INDICATIONS:
Benign gastric and duodenal ulcer, prevention of relapse of duodenal ulcer, Zollinger - Ellison syndrome, Reflux esophagitis, Gastritis and other HCl hypersecretory conditions.

DOSAGE AND ADMINISTRATION:
Doudenal Ulcer, Benign Gastric Ulcer:
Adults & Children above 12 years:
One teaspoonful (5ml) once or twice a day, or as, directed by the physician.
The recommended dose of **Reducid** is 40mg daily taken at bed time. Treatment should be given for 4 to 8 weeks but duration of treatment may be shortened if endoscopy reveals that ulcer is healed.
For prevention of recurrence of doudenal and benign gastric ulcer, the recommended dose is 20mg at bed time given for 1 year.
Zollinger-Ellison Syndrome:
The recommended dose is 20mg 6 hourly. Dosage should be adjusted to the individual patient's need and should continue as long as clinically indicated.
Gastro-Esophagial Reflux Disease:
The recommended dosage for the symptomatic relief of gastro esophageal reflux disease is 20mg **Reducid** twice daily. For the treatment of esophageal erosion or ulceration associated with GERD, the recommended dosage is 40mg **Reducid** twice daily. For the prevention of recurrence